

Ethnobotanical Study of Toxic Plants in Ngadiwono Village, Tosari District, Pasuruan Regency, East Java

Anggraeni Oktavia^{1*}, Serafinah Indriyani², Jati Batoro²

¹Postgraduate Program, University of Brawijaya, Malang, Indonesia

²Department of Biology, Faculty of Mathematics and Natural Sciences, University of Brawijaya, Malang, Indonesia

Abstract

The society in Ngadiwono village is part of Tengger tribe that depends on their surrounding environment on fulfilling the life necessities. However, the society knowledge about toxic plant has never been revealed. Therefore, the main objective of this study is to documenting the toxic plants in Ngadiwono village according to society knowledge and scientific study on its toxic content. This study was conducted in Ngadiwono Village, Tosari District, Pasuruan Regency. The informants were chosen by using snowball method (n=14). Interview was conducted using semi-structural method. The collected data was analysed to obtain ICS value (Index Cultural Significance) and UVs (Use Value). The identification of toxic compound was based on previous study. The study result identified 8 plants that considered to be toxic by local society: *bedor* (*Girardinia palmata* Blume.), yellow *kecubung* (*Brugmansia suaveolens* Bercht. & J.Presl), white *kecubung* (*Brugmansia suaveolens* Bercht. & J.Presl), *jarak* (*Ricinus communis* L.), yellow *terpasan* (*Cestrum elegans* (Brongn.) Schltl), red *terpasan* (*Cestrum elegans* (Brongn.) Schltl), kudisan (*Euphorbia pulcherrima* Willd.), and *ciplukan* (*Physalis peruviana* L.). The highest ICS value was found in *jarak* (*Ricinus communis*). Meanwhile, the lowest ICS value was found in yellow and red *terpasan* (*Cestrum elegans*) due to its minimum use by local society. The highest UVs was found in *kudisan*.

Keywords: Ethnobotany, Ngadiwono, Plant, Tengger, Toxic

INTRODUCTION

The society in Ngadiwono village, part of Tengger tribe, lives in the area of Bromo Tengger Semeru National Park (BTSNP). Most of Tengger society depends on their surrounding environment in occupying their life necessities [1] [2]. Meanwhile, the knowledge of local society is influenced by the local history, custom, and natural resources. In recent years, activities aimed at collecting information, documenting, and confirming the use of ethnomedicine for traditional treatment have been widely conducted in Tengger [2]. Half of the published paper is related with the biological activities of plants, including antimicrobial, molluscicidal, antimalarial, toxicological, and anti-tumor related activities [3]. Previous study mentioned that BTSNP has approximately 118 species of medicinal plants that could be used to medicate 60 types of diseases [4]. However, the knowledge of local society about toxic plants has never been revealed or researched. Therefore, the objective of this

study is to documenting the toxic plant in Ngadiwono village according to society knowledge and scientific study about its content that influenced the toxicity.

MATERIAL AND METHOD

This study was conducted in Ngadiwono Village, Tosari District, Pasuruan Regency, East Java Province, Indonesia (34°20'35.29"E-35°09'27.04"E; 05°19.12"N - 05°53'53.81"N). Ngadiwono village is a buffer zone in Bromo Tengger Semeru National Park that consists of 4 regions: Ledoksari, Krajan, Ketuwon, and Banyu Meneng. Total area of Ngadiwono village is 639.03 hectares with the lowest temperature of 10°C. The distance between settlement area and forest is 2 km. Total male population in BTSNP was 1097, while female was 1474 [5].

Data Collection

Data was collected by using semi-structural and deep interview. The informants were chosen

Correspondence address:

Anggraeni Oktavia

Email : anggraeni_oktavia@yahoo.com

Address : Rajawali no. 14 Sukun Malang East Java

according to snowball method (n=14). We were accompanied by local people during the research, especially when collecting the toxic plants. We terminated the data collection when we already obtained saturated data.

The interview result was analysed to obtain the use value of toxic plant using ICS (Index Cultural Significance). ICS value is an important factor that influencing the presence and use of plants. ICS value was calculated according to the formula listed in the Manual of Flora Diversity Data Collection [6].

$$ICS = \sum_{i=1}^n (q \times i \times e) n_1$$

Where: **ICS** = Index Cultural Significance, is the sum of one individual "use" value from 1 to n; **q** = quality value, is calculated by giving score based on individual plant quality under the criteria: 5 (primary food), 4 (secondary food or primary material), 3 (other food material or medicinal plant), 2 (used in a ritual, mythology, or recreation) and 1 (infrequent use/mere recognition); **i** = intensity value, is the intensity of use under the criteria: 5 (very high intensity), 4 (high intensity), 3 (moderate intensity), 2 (low intensity) and 1 (very low intensity); **e** = exclusivity value, conditions: 2 (the most favored plants that become the main preferences and could not be replaced), 1 (the favored plants but could be replaced by other species of plants) and 0,5 (the beneficial plants that become the secondary preferences).

The obtained data was analysed quantitatively by using UVs equation:

$$UVis = \frac{\sum i Uvis}{nis}$$

Where: Uvs = total number of use that stated by the informant (i) for each plant species (s) in every interview; nis = total number of interview with informant (i) for each plant species (s).

RESULT AND DISCUSSION

The majority of informants were aged over 30 years old with the main occupation as farmer or cattleman (cows or pigs). With respect to the occupation, the informants were predicted to be very accustomed with plant species that could endanger the livestock and considered to be very selective on choosing plants for the feed. They usually use herbicide to inhibit the growth of toxic plant or just cut the plants.

According to the result study, 8 plants were considered to be toxic by the society of Ngadiwono Village (table 1): *bedor* (*Girardinia palmata*), yellow *kecubung* (*Brugmansia suaveolens*), white *kecubung* (*Brugmansia suaveolens*), *jarak* (*Ricinus communis* L), yellow *terpasan* (*Cestrum elegans*), red *terpasan* (*Cestrum elegans*), *kudisan* (*Euphorbia pulcherrima*), *ciplukan* (*Physalis peruviana*).

Bedor was very hard to find in farms. However, this plant still easy to be found near the river, ravine, or forest edge. In addition, *terpasan* (*Cestrum elegans*) also considered to be toxic by local society. There are 3 types of petal on *terpasan*, purple, red or black, and yellow. Interestingly, the purple *terpasan* was used as food for livestock. Meanwhile, the red or yellow *terpasan* was believed to cause poisoning in the livestock with indications of bloated stomach and the appearance of foam from mouth. If they were not treated immediately, they would meet the worst case, mortality.

Kecubung (*Brugmansia suaveolens*) also considered to be toxic by the local society. There are 3 types of *kecubung* in Ngadiwono Village: *kecubung* with yellow petal, *kecubung* with double white petals, while the last is *kecubung* with single white petal. The double white petal *kecubung* was used for feeding the livestock and considered to be non-toxic. We found that local society used water inside the flower buds as eye drops. In contrary, *kecubung* with single white petal, also called as black *kecubung*, and yellow petal were believed to be toxic by the society. The society did not used both *kecubung* for feeding the livestock due to its effect that would cause bloated stomach and puke. In addition, it would cause headache and hallucination if it is consumed by human.

Ciplukan (*Physalis peruviana*) was considered to be toxic by the society in Ngadiwono village. *Ciplukan* would cause bloated stomach for those livestock that consumed it. The side effect was shown by a swelling belly. In contrary, the leaves and fruits of *ciplukan* could be consumed by human and were believed to be a medicine for diabetes.

Kudisan (*Euphorbia pulcherrima*) could be easily found as decorative plants. The exposure of *kudisan* sap would induce itchiness to the skin, therefore, the society classified *kudisan* as toxic plant. The side effect of *kudisan* exposure became the main reason for its local name (*kudis* = mange).

The use of toxic plants by society was calculated using ICS (Index Cultural Significance). The highest ICS value was found in *jarak* (45). The result demonstrated that *jarak* plant was very well known and broadly used by the society. *Jarak* seeds usually used at traditional ceremony called *entas-entas*. In addition, *jarak* seeds contained oil that could be used for fireplace material. *Jarak* seeds also used as a lighting at the burial of the baby umbilical cord. This seeds became an important requirement that must be brought while visiting a shaman. A small quantity of *jarak* seeds (3-5 seeds) could be used as seasoning. The wood also useful as firewood.

Tengger tribe has some unique characteristics compared to other tribe in Java. The majority of Tengger society is centered on the Hindu-Budha religion combined with the traditional customs and beliefs, moreover they still holding tight to their traditional beliefs until now. The society uses plants on their traditional ceremony as *sesajen* (the sacred food) or as symbols of an activity.

Tengger society used *bedor* (ICS = 12, Uvs = 1.077) for a ritual dance called *jaran kepeng*. *Bedor* was struck or given as food for the dancer to assure that they are truly possessed by the spirit, so they become resistant to the itchiness and burning sensation generated by *bedor*. According to the informant, the itchiness could be treated by spreading *ganjan* leaves (*Artemisia vulgaris*) that has been squeezed to the exposed skin.

The stem of white and yellow *kecubung* (ICS = 24, UVs = 1.33) was used to support pipes of an irrigation system in the farm and as firewood. Beside that, *kecubung* that grew near the water resources was believed to maintain the quality and quantity of water.

The leaves and fruits of *ciplukan* (ICS = 9; UVs = 0,835) could be consumed. The society believed that this plant could be used as a medicine for diabetes. *Kudisan* leaves (ICS = 12, Uvs = 1,65) were consumed by the society as *krawu* (mixed vegetable) or as a stuffing in pork belly dish. In contrary, *terpasan* has never been used by the society. The society preferred to prevent the growth of *terpasan* by using herbicide or cut the plants.

The common symptoms of poisoning cases on the livestock were indigestion, such as bloated or swollen stomach. Toxic plants mentioned by the society consist of alkaloids, steroids, and terpenes, that acted as a poison for the stomach. Recent study conducted by Fahrauk et al. [7] stated that LC 50 of ethanol extract on *terpasan*

(*Cestrum nocturnum*) was 44.9658 µg/ml. In fact, this ethanol extract was potentially toxic for *Artemia salina* due to its LC 50 value that was less than 1000 µg/ml. The compound on *terpasan* extract consisted of alkaloids, flavonoid polyphenols, tannins, saponins, monoterpene and sesquiterpenes, also steroids and terpenoids. All of these compounds were proven on interfering the digestion system of shrimp larvae after a series of experiments. These compounds worked by inhibiting the sensory receptors, so the larvae fail to get the stimulus and unable to recognize their food.

The society avoided *jarak* for feeding the livestock due to its effect. A certain amount of *jarak* was able to poisoning human. Anuj [8] discovered the toxic compound of other plant that belong to *jarak* family (*Jatropha curcas*). Latex in seeds contains proteolytic enzymes, curcin and octapeptide, curacycline A, which shows anticomplementary activity. Seed oil contains curcanoleic acid or jatrophin. As well as *Ricinus communis*, *jarak* also contains ricin, such as toxalbumin, phorbol, and cyanic acid, that caused hepatotoxicity in human. The greatest toxic content was found in the seed of *Ricinus communis*. In fact, toxalbumin inhibits the synthesis of 14 proteins, and further effect is gastroenteritis or inflammation on the digestive track. The symptom of toxalbumin poisoning is burning sensations in the throat, stomachache, nausea, vomit, diarrhea, dehydration, lethargy, sleepy, and shock. Other study also mentioned other symptom, such as muscle twitching and an excessive saliva. The toxic effect depends on the doses, whereas the consumption of three seeds could cause a temporary or mild symptom. Moreover, the roasted seed could worsen the effect. Indonesia National Agency of Drug and Food Control [9] described that curcin and ricin would cause a severe gastrointestinal inflammation, characterized by hemorrhage, irritate the gastric mucosa and influence the blood clotting process. The other chemical compound in *jarak* seed is gallic acid, that could cause stomach irritation and kidney failure in a certain amount. First aid for gallic acid poisoning is not to induce vomiting because it could trigger irritation on the gastrointestinal track. For conscious victims, it's better to ask them swallow an activated carbon in less than 4 hours after the exposure. Activated carbon could assist the toxin absorption. The poisoning symptoms generally appear after 1-6 hours after consuming *jarak* seed. Hepatotoxicity occurs 48-71 hours after the toxic material being

consumed. There is no antidotes found for jatropha seed poisoning yet.

Tengger society explained that *kecubung* plant (*Brugmansia suaveolens*) could cause poisoning on the livestock with symptoms bloated and swollen stomach, and then enervation. Moreover, consuming the roasted seed could induce a hallucinatory effect. A study by Yoon (2014) revealed an accident experienced by Koreans. Yoon explained that a Korean used *kecubung* flower as a food decorator on bibimbap (traditional food) and eat a few of it. What happened next was he/she lost consciousness for approximately 10 hours. *Kecubung* contains trophic alkaloids, such as scopolamine, hyoscyamine and atropine. The worst case is all parts of *kecubung* contain toxin, especially in the roots and seeds [10]. Alkaloids in plants, especially scopolamine, cause post-synaptic competitive inhibition of muscarinic cholinergic receptor in center and peripheral areas. *Kecubung* also has anticholinergic effect that appears when poisoning.

The symptoms of *kecubung* poisoning occurs in a short time, approximately 5-10 minutes after consuming *kecubung*. The patients would release an excessive amount of saliva and sweat, followed by dryness of the mouth, mydriasis, and tachycardia. A higher dose would cause urinary retention and hyperthermia. At this point, the patients would experience agitation and anxiety as a result of acute anticholinergics. The patient neuropsychological could be rated by the decreased orientation, chaotic mind, and hallucination. At higher doses, the patient would experience seizures, paralysis, and if not treated properly would cause mortality [11].

In the worst cases, the patient could be administered using Cholinesterase inhibitor physostigmine as an antidote through intravenous route. An activated charcoal (carbo adsorbent) could be administered approximately ≤ 48 hours after consuming *kecubung* [10]. Each flower contains ≈ 0.65 mg of scopolamine and 0.3 mg of atropine. It has been reported that the lethal dose of atropine is 10 mg, so consuming 10 flowers would give a fatal effect [12].

The Ngadiwono society explained that *Ciplukan* (*Physalis peruviana*) could induce bloated stomach on the livestock. However, they added that this plants could be used as a medicine. Layyine [13] demonstrated different LC 50 value that depends on the solvent used in *ciplukan* extract. The solvents were ethanol, n-hexane, and ethyl acetate with LC 50 values of 37,

3, and 496 ppm, respectively. This result demonstrated that ethanol extract has a great potential as antimicrobial, whereas n-hexane has potential as anticancer, and ethyl acetate has potential as a natural pesticide.

The antimicrobial activity on *ciplukan* extract has been investigated by Silva et al. [14]. They revealed that fisalin B, an active compound included in steroid group, plays an important role in antimicrobial activity. Types of fisalin that usually found in *ciplukan* are fisalin B, D, F, and G [15]. In the same concentration of 200 $\mu\text{g/mL}$, fisalin B could inhibit approximately 85% microbes from total microbes that could be inhibited by other fisalin. Fisalin also acts as anti-inflammatory [16] and molluscicides [17]. Fisalin E acts as an anti-inflammatory because of its interaction with glucocorticoid receptors [18].

Ethanol extract (70%) of *ciplukan* is discovered to be toxic for brine shrimp larvae according to its LC 50 that lower than 1000 $\mu\text{g/mL}$, only 39,63 $\mu\text{g/mL}$. If converted, the doses used on the experiment could not exceed 5 gram/kg.bb [18]. Our study demonstrated that *ciplukan* could be useful in a certain dose. More than 5.0 gram/kg.bb *ciplukan* would cause toxicity. The *kudisan* sap contains chemicals called ester, euphorpen, terpenoid, and saponin. The exposure of this compound would cause skin irritation: such as redness, swelling, and itching. The sap exposure on the eye causes mild conjunctivitis. There is no antidotes to treat skin irritation and mild conjunctivitis caused by *kudisan* [19].

CONCLUSION

There were eight plants that considered as toxic by the society of Ngadiwono village. According to the scientific studies, those eight plants have toxic chemicals. Those eight toxic plants were Bedor (*Girardinia palmata*), Kecubung Bunga Kuning (*Brugmansia suaveolens*), Kecubung Bunga Putih (*Brugmansia suaveolens*), Jarak (*Ricinus communis*), Terpasan Kuning (*Cestrum elegans*), Terpasan Merah (*Cestrum elegans*), Kudisan (*Euphorbia pulcherrima*), Ciplukan (*Physalis angulata*). The highest ICS value (45) was found in jarak plant (*Ricinus communis*). While the lowest ICS value was found in yellow and red Terpasan (*Cestrum elegans*) due to its low usage by the society. The highest Uvs value was found in kudisan (1,68).

ACKNOWLEDGEMENT

This section describes gratitude to those who have helped in substance as well as financially.

REFERENCES

- [1]. Batoro, J. 2012. *Etnobiologi Masyarakat Tengger Di Bromo Tengger Semeru Jawa Timur*. Dissertation. Pascasarjana Program. Bogor Agricultural Institute. Bogor.
- [2]. Hakim, L. (2011). Cultural Landscapes of the Tengger Highland, East Java. In *Landscape Ecology in Asian Cultures* (pp. 69-82). Springer Japan.
- [3]. Batoro, J. 2015. *Mengenal Jenis-Jenis Tumbuhan dan Kegiatan Masyarakat Desa Hutan di Kawasan Bromo Tengger Semeru*. Malang: Selaras.
- [3]. Ahmad, I., F. Aqil, and M. Owais. 2006. *Modern Phytomedicine: Turning Medical Plants into Drugs*. Wiley-VCH. Verlag GMBH & CO.KGaA. Germany.
- [4]. Batoro, J. 2015. *Mengenal Jenis-Jenis Tumbuhan dan Kegiatan Masyarakat Desa Hutan di Kawasan Bromo Tengger Semeru*. Selaras. Malang.
- [5]. Anonymous. 1999. *Potret Desa Penyanga Taman Nasional Bromo Tengger Semeru*. Departemen Kehutanan (Bagian Proyek Pemantapan Pengelolaan Taman Nasional Bromo Tengger Semeru Tahun Anggaran 1998/1999). Malang.
- [6]. Hakim, L. (2014). *Etnobotani dan Manajemen Kebun-Pekarangan Rumah: Ketahanan Pangan, Kesehatan dan Agrowisata*. Selaras. Malang.
- [7]. Fahrak, F., R. Julia, and N. Neng. 2014. Uji Bioaktivitas Ekstrak Daun dan Fraksi Daun Kembang Dayang (*Cestrum nocturnum* Linn.) Terhadap *Artemisia salina* Leach dengan menggunakan Metode BSLT (Brine Shrimp Lethality Test). *Proceeding Seminar Nasional Ilmu Pengetahuan dan Teknologi Universitas Jenderal Achmad Yani*. 52-53.
- [8]. Gupta A., K. Alok, A. Ajay, Motoki, and V. Archana. 2016. Acute accidental mass poisoning by *Jatropha curcas* in Agra, North India, Egypt. *J Forensic Sci* <http://dx.doi.org/10.1016/j.ejfs.2016.04.002>
- [9]. BPOM. 2015. Keracunan Biji Jarak. accessed on 1 february 2017 <http://ik.pom.go.id/v2015/qa/keracunan-biji-jarak> diakses
- [10]. Gopel C, Laufer C, Marcus A. 2002. Three cases of angel's trumpet tea-induced psychosis in adolescent substance abusers. *Nord J Psychiatry*. 56:49–52
- [11]. Hall RC, Popkin MK, McHenry LE. 1977. Angel's Trumpet psychosis: a central nervous system anticholinergic syndrome. *Am J Psychiatry*. 134:312–314.
- [12]. Greene GS, Patterson SG, Warner E. 1996.: Ingestion of angel's trumpet: an increasingly common source of toxicity. *South Med J* 1996, 89:365–369.
- [13]. Layyina, H. 2014. Toksisitas Ekstrak Ciplukan (*Physalis angulata*) Berdasarkan Uji Letalitas Larva Ugang. Departemen Kimia. Fakultas Matematika Dan Ilmu Pengetahuan Alam. Institut Pertanian Bogor.
- [14]. Silva MTG, Simas SM, Batista TGFM, Cardarelli P, Tomassini TCB. 2005. Studies on antimicrobial activity, in vitro, of *Physalis angulata* L. (Solanaceae) fraction and physalin B bringing out the importance of assay determination. *Mem Inst Oswaldo Cruz*. 100:779-782. Silve et al., 2015
- [15]. Sa, M.S., M.N. Menezes, A.U. Krettli, I.M. Ribeiro, T.C.B. Tomassini, R.R. Santos, W.F. Azevedo, M.B.P. Soares. 2011. Antimalarial activity of physalins B, D, F, and G. *J. Nat. Prod*. 74. 2269-2272.
- [16]. Pinto, N.B., T.C. Morais, K.M.B. Carvalho, C.R. Silva, G.M. Andrade, G.A.C. Brito, M.L. Veras, O.D.L. Pessoa, V.S. Rao, and F.A. Santos. 2010. Topical anti-inflammatory potential of physalin E from *Physalis angulata* on experimental dermatitis in mice. *Phytomedicine*. 17. 740-743.
- [17]. Santos, J.A.A., T.C.B. Tomassini, D.C.D. Xavier, I.M. Ribeiro, M.T.G. Silva, and Z.B.M. Filho. 2003. Molluscicidal activity of *Physalis angulata* L. extracts and fractions on *Biomphalaria tenagophila* (d'Orbigny, 1835) under laboratory conditions. *Mem Inst Oswaldo Cruz*. 98. 425-428.
- [18]. Baedowi. 1998. Timbunan Glikogen dalam Hepatosit dan Kegiatan Sel Beta Insula Pancreatis Tikus Putih (*Rattus norvegicus*) Akibat Pemberian Ekstrak Daun Ciplukan, Penelitian Tanaman Obat di Beberapa Perguruan Tinggi di Indonesia IX. Departemen Kesehatan RI. Jakarta.
- [19]. The American Society For Prevention of Cruelty to Animal. Toxic and Non Toxic Plant. 2017. <https://www.aspc.org/pet-care/animal-poison-control/toxic-and-non-toxic-plants/poinsettia> Anonim, 2017?

Table 1. List of toxic plants and the exposure effect.

No.	Local Name	Scientific Name	Family	Toxic Part	The exposure side effect
1.	Bedor	<i>Girardinia palmata</i>	Urticaceae	Thorn on the leaves and branches	Skin burning sensation
2.	Red Terpasan	<i>Cestrum elegans</i>	Solanaceae	Branches, leaves, and flowers	Bloated stomach and mortality on livestock
3.	Yellow Terpasan	<i>Cestrum elegans</i>	Solanaceae	Branches, leaves, and flowers	Bloated stomach and mortality on livestock
4.	Jarak	<i>Ricinus communis</i>	Euphorbiaceae	Leaves and seeds	Bloated stomach on livestock, seeds could cause hallucination on human
5.	Kecubung with white single petal/black kecubung	<i>Brugmansia Suaveolens</i>	Solanaceae	Leaves and seeds	Bloated stomach on livestock, seeds could cause hallucination on human
6.	Yellow Kecubung	<i>Brugmansia Suaveolens</i>	Solanaceae	Leaves and seeds	Bloated stomach on livestock, seeds could cause hallucination on human
7.	Ciplukan	<i>Physalis peruviana</i>	Solanaceae	Leaves	Bloated stomach on livestock
8.	Kudisan	<i>Euphorbia pulcherrima</i>	Euphorbiaceae	Sap from cut branches	Itchiness on the parts that exposed by sap

Table 2. ICS value according to the benefit of toxic plant by the society of Ngadiwono Village.

Name	ICS	UVs	Benefit	Usage Procedure
Bedor (<i>Girardinia palmata</i>)	12	1.077	<i>Jaran kepong</i> ritual dance	All plant part is struck to the dancer body.
Red Terpasan (<i>Cestrum elegans</i>)	0.5	0		Not used
Yellow Terpasan (<i>Cestrum elegans</i>)	0.5	0		Not used
Jarak (<i>Ricinus communis</i>)	45	1.27		
Seed			Seasoning Ritual lighting	2-5 seeds One seed is stuck to a bamboo and burned
Stem			Firewood	
Leaf			Fruit wrapper	Wrap banana or raw fruit to accelerate the ripening process
Fruit			Medicine for sprue	
Black Kecubung (<i>Brugmansia suaveolens</i>)	24	1.33		
Stem			Firewood The guardian of water resources	Planted near the water resources
Yellow Kecubung (<i>Brugmansia suaveolens</i>)	24	1.33	The guardian of water resources	Planted near the water resources
Stem			Firewood	
Ciplukan (<i>Physalis peruviana</i>)	9	0.835	Traditional medicine	A handful leaf is boiled, then drunk
Kembang Kudis (<i>Euphorbia pulcherrima</i>)	24	1.68	Fever medicine	A handful leaf is boiled, then drunk
Leaf			Vegetable	Put inside the pig belly and roasted or used as the ingredient of <i>krawu</i>
			Decorative plant	Planted in front of the house